



Annual Review of Plant Biology

Evolution of Plant NLRs: From Natural History to Precise Modifications

Janina Tamborski and Ksenia V. Krasileva

Department of Plant and Microbial Biology, University of California, Berkeley,
California 94720, USA; email: kseniak@berkeley.edu

Annu. Rev. Plant Biol. 2020. 71:11.1–11.24

The *Annual Review of Plant Biology* is online at
plant.annualreviews.org

<https://doi.org/10.1146/annurev-arplant-081519-035901>

Copyright © 2020 by Annual Reviews.
All rights reserved

Keywords

NLRs, evolution, protein engineering, plant immunity

Abstract

Nucleotide-binding leucine-rich repeat receptors (NLRs) monitor the plant intracellular environment for signs of pathogen infection. Several mechanisms of NLR-mediated immunity arose independently across multiple species. These include the functional specialization of NLRs into sensors and helpers, the independent emergence of direct and indirect recognition within NLR subfamilies, the regulation of NLRs by small RNAs, and the formation of NLR networks. Understanding the evolutionary history of NLRs can shed light on both the origin of pathogen recognition and the common constraints on the plant immune system. Attempts to engineer disease resistance have been sparse and rarely informed by evolutionary knowledge. In this review, we discuss the evolution of NLRs, give an overview of previous engineering attempts, and propose how to use evolutionary knowledge to advance future research in the generation of novel disease-recognition capabilities.



Contents

1. INTRODUCTION	11.2
1.1. Extracellular and Intracellular Branches of Plant Immune Perception	11.2
1.2. NLRs Are Intracellular Immune Receptors	11.3
1.3. Mechanisms of NLR Activation	11.3
2. GENOMIC BASIS OF NLR EVOLUTION	11.6
2.1. NLR Evolution Across Angiosperms	11.6
2.2. Lineage-Specific Clade Expansions and Contractions	11.7
2.3. NLR Allelic Diversity and Novel Gene Fusions	11.7
2.4. Functional Networks Among Sensor and Helper NLRs	11.8
3. PATTERNS OF CONVERGENT EVOLUTION IN NLR BIOLOGY	11.8
3.1. Dependence of Sensors on Helpers Arose More Than Once During NLR Evolution	11.8
3.2. Regulation of NLRs by Small RNAs	11.10
3.3. Similarities to Mammalian Intracellular Immunity	11.10
4. CURRENT PROGRESS IN ENGINEERING NLR ALLELES	11.13
4.1. NLR Chimeras Define Effector Binding Regions and Allow Transfer of Recognition Between Allelic NLRs	11.13
4.2. Introduction of New Effector Substrate Sequences into Plant Proteins	11.13
4.3. Introduction of Amino Acid Changes to Modify Effector Recognition Specificity	11.13
5. PROSPECTS AND CHALLENGES IN THE ENGINEERING OF NLRs	11.14
5.1. Engineering Guardees for NLR Activation upon a Wide Range of Effector Activities	11.14
5.2. Engineering NLRs with New Integrated Domains	11.15
5.3. Exploiting NLRs with Naturally Diverse LRR Binding Specificities	11.15
6. CONCLUSIONS AND FUTURE DIRECTIONS	11.16

1. INTRODUCTION

1.1. Extracellular and Intracellular Branches of Plant Immune Perception

Plant immune receptors monitor the extracellular and intracellular compartments for signs of a pathogen invasion. Numerous plasma membrane-localized receptors, commonly referred to as pattern recognition receptors, surveil the extracellular space for signs of an invading pathogen. They recognize conserved molecular patterns derived from microbial molecules (microbe-associated molecular patterns) or from the actions of microbial molecules (damage-associated molecular patterns), collectively referred to as danger signals (57). The recognition of danger signals leads to the activation of well-characterized immune responses that include but are not limited to the production of reactive oxygen species (ROS), an increase in cytosolic calcium concentration, activation of mitogen-activated protein (MAP) kinase cascades, and the expression of defense genes (18). Pattern recognition receptors, their signaling partners, and their outputs have been reviewed elsewhere (37, 75).

Cytoplasmic immune receptors recognize foreign molecules that are secreted into the plant cell by an invading pathogen as well as enzymatic modifications to plant components. These are called nucleotide-binding leucine-rich repeat receptors (NLRs) after the two core domains shared

Nucleotide-binding leucine-rich repeat receptors (NLRs): intracellular immune receptors in plants, animals, and fungi

11.2 Tamborski • Krasileva



Review in Advance first posted on
February 24, 2020. (Changes may
still occur before final publication.)

among the majority of these proteins. The secreted pathogen molecules are termed effectors, and their function supports invasion of the pathogen, often through the suppression of plant immunity. The recognition of effectors by NLRs leads to some of the same responses as the recognition of extracellular patterns. These include the accumulation of ROS, activation of MAP kinase cascades, and defense gene expression; however, the kinetics of these responses differs from that of surface receptors (45). In addition, NLR activation culminates in localized cell death, known as the hypersensitive response (HR).

1.2. NLRs Are Intracellular Immune Receptors

NLRs have a distinct domain architecture that consists of a nucleotide-binding (NB-ARC) domain and a series of C-terminal leucine-rich repeats (LRRs), and most have an N-terminal extension consisting of a Toll/interleukin-1 receptor (TIR) domain, a coiled-coil domain (CC), or a divergent coiled-coil domain (CC_R) similar to the RESISTANCE TO POWDERY MILDEW 8 (RPW8) domain (113). NLRs form three classes on the basis of the N-terminal domain and evolutionary history of the NB-ARC: TIR-NLRs (TNLs), CC-NLRs (CNLs), and RPW8-NLRs (RNLs) (**Figure 1a**) (113). A few NLRs do not have all of the canonical domains; for instance, RESPONSE TO THE BACTERIAL TYPE III EFFECTOR PROTEIN HOPBA1 (RBA1) is a TIR-only protein that recognizes the bacterial effector HopBA1 (96), and proteins containing only TIR or RPW8 domains can elicit resistance to pathogens on their own (94, 135, 144). NLRs can directly bind and recognize effectors (e.g., 46, 62) or indirectly recognize the modification of another plant component through effector function (129, 150). Plants also have immune receptors with integrated domains (IDs) that mimic pathogen targets and are activated in response to modification by the effectors (28, 70, 105).

In general, we can divide NLRs into two functional groups: direct/indirect sensor NLRs that are involved in the recognition of invasion and helper NLRs that are genetically required by other NLRs for immune activation (65). In some cases, NLRs work in a pair consisting of a sensor and a helper and are genetically linked. In some species, helper NLRs are proposed to have arisen from a single pair before expanding into complicated networks of many NLRs, such as the *NLR REQUIRED FOR CELL DEATH* (*NRC*) gene family, which has expanded in the asterids and serves as helpers for a number of sensor CNLs (140–142). RNL helpers that function downstream of sensor NLRs and are required for both resistance and cell death include ACTIVATED DISEASE RESISTANCE 1 (*ADR1*) and N REQUIREMENT GENE 1 (*NRG1*) (25, 71, 102, 143).

1.3. Mechanisms of NLR Activation

The molecular mechanisms underlying activation of sensor NLRs and of their helpers are expected to differ significantly and are not yet fully understood. However, their shared domains and common evolutionary origin suggest that multimerization through the NB-ARC domain following exchange of ADP for ATP is a key step in NLR activation. For this reason, NLRs are often referred to as molecular switches in immune signaling (123). The ADP-bound state is thought of as the off state, in which the LRR associates with the NB-ARC domain, thereby stabilizing the NLR in the inactive state (124). The activation of NLRs is generally associated with the ATP-bound state and is referred to as the on state.

Cryo-electron microscopy structures of an indirect sensor protein, *Arabidopsis thaliana* HOPZ-ACTIVATED RESISTANCE 1 (*ZAR1*), have demonstrated both states and suggested the existence of a third, intermediate state (133, 134). These structures showed that the ADP-bound form was monomeric and had multiple intramolecular contact points between the LRR and NB-ARC

Effector: pathogen-derived molecule translocated into the plant cell to modify its environment for the benefit of the pathogen

NB-ARC: nucleotide-binding domain of NLRs that acts as a molecular switch and facilitates oligomerization of the active receptor complex

Leucine-rich repeats (LRRs): domains that form a horseshoe structure that facilitates protein–protein interactions

Toll/interleukin-like receptor (TIR) domain: found on its own or as part of NLRs; can induce HR on its own

Coiled-coil (CC) helix bundle: facilitates formation of the signaling platform; can induce HR on its own


RESISTANCE TO POWDERY MILDEW 8 (RPW8): found on its own or as part of NLRs; similar to fungal pore-inducing toxins

TNL: NLR with an N-terminal TIR signaling domain


CNL: NLR with an N-terminal CC signaling domain



a Phylogenetic relationship of NLRs across monocots and dicots



b NLR functions



(Caption appears on following page)



Figure 1 (Figure appears on preceding page)

NLR evolution and functions in flowering plants. (a) Maximum likelihood phylogeny of 7,133 NLRs from 11 dicots (green), 7 monocots (yellow), and 1 moss (red) based on NB-ARC domain alignment (11). Major classes of NLRs are depicted as arcs: RNLs (light purple), TNLs (medium purple), and CNLs (dark purple). Examples of well-characterized NLRs with different functions are marked on the tree in blue text (sensors) and gray text (helpers). The tree is rooted on the longest internal branch, and is based on the NB-ARC domain. Bootstrap values >80 are indicated on the tree as black circles. Monocots (Poales) are *Oryza sativa*, *Sorghum bicolor*, *Triticum aestivum*, *Setaria italica*, *Hordeum vulgare*, *Zea mays*, *Brachypodium distachyon*. Dicots are *Arabidopsis thaliana*, *Medicago truncatula*, *Solanum tuberosum*, *Solanum lycopersicum*, *Vitis vinifera*, *Glycine max*, *Malus domestica*, *Prunus persica*, *Eucalyptus grandis*, *Fragaria vesca*, *Populus trichocarpa*. Moss is *Physcomitrella patens*. An interactive version of this tree constructed by Dr. Paul Bailey following the methodology described in Bailey et al. (10), is available at <https://itol.embl.de/tree/149155221225191791507546488>. (b) Schematic diagram of NLR functions. Blue proteins indicate sensors, including direct binders, indirect binders, and NLRs with integrated domains; gray proteins are helpers. Pathogen-derived effectors are shown as circles; plant proteins targeted by the pathogen, as well as integrated domains, are shown as hexagons, and the effector recognition site is a triangle.

[helical domain 1 (HD1) and winged helix domain (WHD)] (134). The recognition of effector-induced changes in guard cells through the LRR alters the protein's conformation, thereby releasing the negative inhibition. The addition of ATP induced an oligomeric state and the formation of a wheel-shaped pentamer called the resistosome (133), reminiscent of the mammalian apoptosome (151) that represents the active state of Apaf-1 and the inflammasomes that assemble upon activation of mammalian NLRs (126). The CC of ZAR1 contributes to its oligomerization by forming an α -helical barrel (133). The active complex associates with the plasma membrane, and the charged residues inside the funnel are required for the initiation of cell death and disease resistance (133).

How ligand binding or modification affects the nucleotide state of the NB-ARC domain needs further elucidation. One possibility, based on observations from numerous studies, is that the binding of an effector leads to conformational changes within the NLR molecule, enabling exposure of the NB-ARC domain and thereby facilitating the exchange from ADP to ATP (86, 125, 137). This is also the case in ZAR1, in which binding of the uridylated host protein kinase PBS1-LIKE 2 (PBL2) to the ZAR1/RKS1 (RESISTANCE-RELATED KINASE 1) complex resulted in the rotation in the NB-ARC relative to the other domains and a loss of affinity to ADP (134). However, in a different model, known as the equilibrium-based switch model, a pathogen effector showed no significant binding affinity to the ADP-bound state of the NLR (15). In this model, the NLRs constitutively cycle between the on and off states, and the pathogen effector stabilizes the on state and shifts the equilibrium in favor of the on state, leading to activation of defense responses (15). Both models probably coexist and are utilized by different NLRs. Allowing different modes of activation would not only make immune responses more robust against pathogen interference but would also allow adaptation to varying effector binding affinities.

To date, no structures of active TNL complexes have been reported; however, it is logical to think that they would assume a similar wheel-shaped structure based on oligomerization of the NB-ARC domain. TNLs are known to form oligomers, and TIR domains can self-associate across multiple surfaces (16, 138, 148). While the active ZAR1 resistosome associates with a membrane through its N-terminal α -helix, the active TNL complex likely signals through the enzymatic breakdown of NAD⁺ by the TIR domains (59, 131). Multiple plant TIRs, as well as a mammalian TIR from STERILE ALPHA AND TIR MOTIF CONTAINING 1 (SARM1), were capable of cleaving NAD⁺ into nicotinamide and cyclic ADP-ribose (cADPR) (59, 131); however, some plant TIRs produced a third product, v-cADPR (131). The enzymatic activity in both systems required self-association and depended on the conserved catalytic glutamate (59, 131).

Activation of NLRs induces downstream responses that can ultimately lead to HR, a form of localized cell death. While all TNLs to date have been found to signal through and strictly require the *ENHANCED DISEASE SUSCEPTIBILITY 1* (*EDS1*) and *PHYTOALEXIN DEFICIENT 4*

RNL: NLR with an N terminus homologous to the RPW8 signaling domain

Integrated domain (ID): a noncanonical domain fused to an NLR that does not share its evolutionary history

Sensor: NLR responsible for binding an effector or recognizing its activity

Helper: NLR that is activated by another NLR or a signaling cascade downstream of effector recognition

Resistosome: a wheel-shaped oligomeric structure comprising NLRs that is assembled upon activation



(*PAD4*) genes encoding for lipase-like proteins (50), exactly how the enzymatic products of the TIR domains affect downstream signaling components remains unknown. Recent studies have substantially added to our understanding of how RNLs function downstream of sensor NLRs (25, 71, 102, 143). The new data suggest that NRG1 interacts with EDS1 and *SENESCENCE-ASSOCIATED GENE 101 (SAG101)* and that this complex is strictly required for initiation of HR in *A. thaliana* (102). This complex might directly execute HR, as the RPW8 domain of NRG1 is reminiscent of pore-forming toxins and it may insert into the plasma membrane to disrupt membrane integrity (11). Some CNLs genetically rely on NON-RACE SPECIFIC DISEASE RESISTANCE 1 (NDR1) (1, 27), a protein that is anchored in the plasma membrane. Since the active resistosome of ZAR1 was associated with membranes, it is possible that NDR1 might be required for the suggested pore-forming activity. Together, these studies have brought us much closer to understanding the initiation and function of HR, which is a crucial step toward a better understanding of NLR-mediated immunity.

In the following sections, we discuss different evolutionary processes that yield the wide variety of NLRs, illustrate how phylogenetic and evolutionary analyses can indicate which modes of activation are deployed by which NLRs, and explain how we can apply this knowledge to engineer novel disease resistance.

2. GENOMIC BASIS OF NLR EVOLUTION

2.1. NLR Evolution Across Angiosperms

From analyses of more than 100 sequenced plant genomes, common patterns of NLR evolution have emerged that were not apparent in studies of a single species or plant family (54, 113, 149). On the basis of their NB-ARC phylogeny, the TNLs, CNLs, and RNLs form three monophyletic groups and have unique N-terminal domains that likely represent ancestral fusions of the TIR domain, CC, and CC_R to an ancestral NB-ARC domain (**Figure 1a**). Representatives of all three NLR types are present in the basal plant lineage *Amborella*, suggesting that the split into the clades is ancient (113). Ancestral reconstruction of NLRs from 22 representative angiosperms suggested that a basal plant had approximately 23 groups of NLRs (90, 113). An orthogroup analysis of 20,571 NLRs across 75 plant genomes demonstrated that only 38 of 311 NLR families are conserved across monocots and dicots, while the other groups represent lineage-specific gene expansions (149).

The copy number of RNLs, TNLs, and CNLs varies across plants (**Figure 1a**). The RNL clade is usually characterized by a low copy number (54, 70, 113, 149), with the exception of gymnosperms (there are 31 ADR homologs in spruce) (145). RNLs show remarkable intron conservation, with *Amborella* and dicots sharing four introns and monocots three introns (the second intron is lost) (90, 113). The separation of NRG and ADR occurred prior to the divergence of angiosperms, and they are still conserved in syntenic blocks across flowering plants (113). The NRG genes, but not the ADR genes, have been lost in several lineages (113).

The TNLs form two subfamilies, TIR1 and TIR2 (105), with only TIR2 NLRs retained in monocots (94, 105). TIR1 NLRs have proliferated in many dicot species but are absent in some dicot lineages (113). Across all flowering plants, TNLs show remarkable conservation of intron/exon junctions: The first intron separates TIR from NB-ARC, the second separates NB-ARC from LRRs, and the third separates the first LRR from the rest of the protein (90, 113). In contrast, CNLs do not share conserved introns (90, 113), and current data suggest that ancestral CNLs were intronless and that introns in CNLs were likely gained later in evolution (113).

The CNLs are subdivided into at least four distinct groups based on amino acid motifs conserved across a large evolutionary distance (139). Two groups shared across monocots and dicots



contain an EDVID motif, such as ZAR1 and RPM1 proteins in *Arabidopsis*, Rx in potato, and Sr33/MLA in Poaceae (139). Some CNLs also contain a functionally conserved N-terminal methionine, alanine, aspartate, alanine (MADA) motif (2) within the first α -helix, which is rearranged upon oligomerization and mediates membrane anchoring (133).

In the context of the global evolutionary history of NLRs (**Figure 1a**), it is evident that sensor and helper roles evolved multiple times. The NRC helper clade emerged before the split of Caryophyllales and asterids more than 100 million years ago and expanded in Solanaceae (140); yet the RNL helper clade is even older, dating to before the split of gymnosperms and angiosperms (113). Similarly, paired NLRs such as *RESISTANT TO RALSTONIA SOLANACEARUM 1/RESISTANT TO PSEUDOMONAS SYRINGAE 4 (RRS1/RPS4)* and *R-GENE ANALOG 4/R-GENE ANALOG 5 (RGA4/RGA5)* are paraphyletic (**Figure 1a**). The sensors *RESISTANT TO PSEUDOMONAS SYRINGAE 2 (RPS2)* and *RESISTANCE TO PSEUDOMONAS SYRINGAE PV. MACULICOLA 1 (RPM1)* guard the same plant protein, *RPM1-INTERACTING PROTEIN 4 (RIN4)*, but diverged at least before the split of monocots and dicots, suggesting that their function likely arose independently. Similarly, genes encoding direct binders that recognize the same pathogen [such as *RECOGNITION OF PERONOSPORA PARASITICA 13 (RPP13)* and *RPP1*] do not cluster together. This finding demonstrates that functional similarity between NLRs does not necessarily reflect their evolution.

2.2. Lineage-Specific Clade Expansions and Contractions

There is a 100-fold variation in the number of NLRs across plant genomes, ranging from a few dozen in papaya, kiwi, cucumber, and watermelon (8, 54, 78, 81, 149) to several thousand in hexaploid wheat (4). Even closely related species can show lineage-specific expansions and contractions (4, 8, 9, 55, 56, 63, 78, 82, 118, 149). While the selection mechanisms driving NLR expansions and contractions remain elusive, they can reflect plant lifestyle and be shaped by the selection pressures from the environment. Plant lifestyles that have been correlated with NLR history include aquatic environment (9) and either being dioecious (kiwi, papaya) or having separate male and female flowers (maize, cucumber) (8). Population genetics of NLR repertoires indicate that environmental pressure shapes NLR diversity within a single species, such as adaptation to pathogens in wild tomato (119, 120) and *Arabidopsis* (128), and can lead to long-term maintenance of polymorphisms (66).

How can the expansion or contraction of NLRs be achieved in a short evolutionary time? A burst of CNL expansions in Solanaceae and Poaceae is attributable in part to the activity of long terminal repeat (LTR) retrotransposons (67). The repetitive nature of NLRs themselves also aids in their evolution, deploying recombination for gene conversions (92, 111) and replication machinery for local duplications (72).

2.3. NLR Allelic Diversity and Novel Gene Fusions

Expansion and contraction of NLR genes can be observed even among closely related species; however, gene duplications alone are not sufficient to generate new pathogen recognition. The rapid NLR evolution proposed in the so-called birth-and-death model (91) holds true today. Current genomic data sets support the diversification of NLRs through intragenic and intergenic recombination and gene conversion that generate chimeric LRRs (111), as well as point mutations in surface-exposed regions of LRRs (83, 87). Such NLRs are capable of recognizing highly variable and even structurally unrelated effectors, most likely through a direct binding mechanism, and can provide resistance against multiple pathogens (17, 34, 83, 117). NLRs with high allelic variation can arise in either TNL or CNL clades and are not monophyletic.



Guardee: a plant protein guarded by an NLR for signs of pathogen-derived effector modifications

More recently, NLRs with IDs emerged as sources of new recognition specificities (28, 70, 105). A specialist NLR-ID clade in Poaceae, major integration clade 1 (MIC1), continuously shuffles IDs, thereby facilitating the generation of new ID sensors (10, 121). NLR-IDs show evidence of duplications and intrachromosomal translocations, but the exact genomic mechanism of domain integration remains elusive. An analysis of NLR-IDs in wheat suggests that once the ID is integrated near the NLR in the genome, alternative splicing gives rise to a fused transcript (4).

Rapid generation of new recognition specificity can come at a cost: autoimmunity (19, 32). Several allelic variants of NLRs have been linked to autoimmunity in *Arabidopsis* (3, 31, 77), representing a possible mismatch of NLRs and their guardees. An allelic variant of the *A. thaliana* NLR *RPP7* was recently shown to cause an autoimmune response when combined with incompatible alleles of *RPW8* (67). One might speculate that the low number of NLRs in genomes of highly inbred crops such as maize is the result of autoimmunity leading to the loss of mismatched NLRs (8, 118).

2.4. Functional Networks Among Sensor and Helper NLRs

Flor's (51) original gene-for-gene postulate in *R* gene-effector recognition evolved into a new paradigm of functional networks among both sensors and helpers. RPS2 and RPM1 are classic examples of sensor NLRs that can track multiple effectors by guarding the effector target hub RIN4 (7, 13, 39, 84, 85). Other NLRs such as ZAR1 monitor multiple paralogous guardees, each tracking a distinct effector (12, 73, 79, 108, 112, 132). Alleles of the *RRS1* encoding NLR-ID with an integrated WRKY domain can detect at least two effectors, AvrRps4 and PopP2 (42, 89, 106, 107), and the *RRS1/RPS4* pair can provide resistance against three different pathogens (95). While multiple effector recognition specificities can be encoded in one NLR, independently evolved NLRs can also provide recognition of the same effector. In soybean and *Arabidopsis*, RIN4 is guarded by the evolutionarily unrelated NLRs Rpg1b/Rpg1r and RPM1/RPS2, respectively, suggesting that recognition of the *Pseudomonas* effectors AvrRpm1 and AvrB evolved independently (6). Recognition of protease activity of another *Pseudomonas* effector, AvrPphB, evolved at least twice in *Arabidopsis* and barley through NLRs guarding the same target, AVRPPHB SUSCEPTIBLE 1 (PBS1) (24). Finally, orthologous NLRs, such as *MLA/Sr33/Sr50* and *Rx1/Gpa2*, evolved recognition of distinct effectors through allelic variation (83, 117). Altogether, these studies demonstrate that recognition specificity of a sensor NLR cannot be assigned on the basis of sequence identity alone.


Helper NLRs can form functional networks of their own. In *Arabidopsis*, TNLs and CNLs can signal through proteins belonging to the NRG1 and ADR1 subclades, with TNLs showing a stronger preference for NRGs and CNLs for ADRs (25, 71, 102, 143). The NRC helper clade represents a more recent network, supporting a closely related CNL sensor clade (140). Whereas the protein products of paired NLR genes (such as *RG4/RGA5* and *RRS1/RPS4*) also show physical interaction (29, 61), functional dependence of NLRs on NRC and RNL has been shown so far only genetically (25, 71, 102, 140–143). Whether protein-protein interactions are unique to paired NLRs and have been lost during formation of helper networks remains to be determined.

3. PATTERNS OF CONVERGENT EVOLUTION IN NLR BIOLOGY

3.1. Dependence of Sensors on Helpers Arose More Than Once During NLR Evolution

The RNL clade is an ancient, conserved group of helpers shared between monocots and dicots, while the helper function of the NRCs arose independently (**Figure 2a**). The NRC clade most



**Figure 2**

Patterns of convergent evolution in NLR biology. (a) Sensor/helper networks and pairs have evolved independently multiple times during NLR evolution. Sensor NLRs are involved in recognition of pathogen invasion, whereas helpers are genetically required by sensors to execute immune signaling. They interact in networks, as is the case for helpers of the RNL and NRC classes (i), or in a paired relationship (ii), which is usually associated with genetic linkage between the helper and sensor in a head-to-head orientation. Representative examples of each NLR type are shown in parentheses. (b) Regulation of NLRs through small RNAs (sRNAs) is conserved across most plant lineages. The NLRs are regulated by sRNAs, including microRNAs (miRNAs) that are derived from NLRs themselves and involved in transcriptional silencing (i) and sRNAs that target transposable elements (TEs) and NLRs through RNA-dependent DNA methylation (ii). A representative example of such a miRNA is shown in parentheses. (c) The formation of a multimeric complex of activated NLRs is conserved across kingdoms. (i) The known resistosome, as demonstrated by Wang et al. (132, 133). We propose several hypothetical ones (ii, iii) that could be analogous to the variety of inflammasomes formed by mammalian NLRs.

likely emerged from a solanaceous NLR pair more than 100 million years ago and has since evolved into a functional network of helpers and sensors (140). Even more recently, two independent clades of TNLs and CNLs, exemplified by *RRS1/RPS4* and *RGA5/RGA4*, subdivided into sensors and helpers in both Poaceae and Brassicaceae (28, 95). Intriguingly, Poaceae and Brassicaceae genes encoding sensor and helper pairs are genetically linked in head-to-head orientation, and the sensor often carries an ID (Figure 2b) (10, 95). The genomic and evolutionary mechanisms that drove convergent head-to-head orientation of gene pairs, fusion with IDs, or functional subdivision into helpers and sensors are currently unknown.



The dependence of sensor immune receptors on additional receptors is not unique to plants. In animals, the NLR FAMILY, APOPTOSIS INHIBITORY PROTEIN 5 (NAIP5) is a flagellin-binding sensor that depends on the helper NLR FAMILY, CASPASE ACTIVATION AND RECRUITMENT DOMAIN-CONTAINING 4 (NLRC4) to execute the response (126). Even outside the NLR family, ligand-binding receptor kinases such as FLAGELLIN SENSING 2 (FLS2) and BRASSINOSTEROID INSENSITIVE 1 (BRI1) that functionally act as sensors depend on the physical interaction with the coreceptor BRI1-ASSOCIATED RECEPTOR KINASE 1 (BAK1) for complex activation and downstream signaling (35, 93). Therefore, the requirement of an additional partner by ligand-sensing receptors is a general evolutionary theme. It is important to understand whether such a requirement is imposed by common functional constraints for receptors to initiate signaling cascade or whether it presents an additional level of regulation.

3.2. Regulation of NLRs by Small RNAs

Misregulation of NLRs is costly; therefore, they are tightly regulated on both the messenger RNA and protein levels (76, 115). A global analysis of NLRs in plant genomes revealed a common pattern: NLRs are regulated by microRNAs (miRNAs) that can be generated from duplicated NLRs (**Figure 2b**). Some miRNAs, such as miR482/2118, which targets CNLs, appeared a long time ago; these miRNA families are conserved across most plant lineages and target the conserved P-loop region of the NB-ARC domain (54). Other miRNAs appeared more recently and are lineage specific (58, 76, 80, 99, 110, 115, 145, 147, 149). Continuous evolution of new miRNAs has been linked to lineage-specific expansion of NLRs, with new miRNAs likely to be derived from inverted duplications of target NLR sequences (149).

Regulation of NLRs by miRNAs bears similarity to another group of rapidly proliferating genetic elements that are also regulated by small RNAs (sRNAs): transposable elements (TEs). Both NLRs and TEs duplicate rapidly, show lineage-specific expansions and contractions, and can jump around in the genome. Both newly inserted TEs and recently duplicated genes are regulated by epigenetic marks through RNA-dependent DNA methylation guided by sRNAs (97, 103). A global analysis of sRNAs and associated changes in *A. thaliana* epigenetic marks upon *Pseudomonas syringae* infection showed that both TEs and NLRs are derepressed early in infection and start to produce sRNAs, many of which map to both NLR and TE loci (23). While TEs are silenced later in the course of infection by the process of RNA-directed DNA methylation (RdDM) depositing de novo DNA methylation at the genomic loci, many NLRs, including ADR1, continue to be expressed, despite the continuous presence of sRNAs that can target them (23).

Coregulation of NLRs by TEs inserted into the promoter elements has an important functional role (**Figure 2b**). In rice, the insertion of a miniature inverted repeat transposable element into the promoter of the NLR gene *PigmS* constrains its expression to pollen and silences its transcription through RdDM in other plant tissues (41). Functionally, the *PigmS* protein is a dominant suppressor of another NLR, *PigmR*, which confers resistance to *Magnaporthe oryzae*. When *PigmS* is silenced, the *PigmR* protein is active and provides resistance in vegetative tissues; however, in pollen, it is suppressed by *PigmS*, thereby reducing the yield penalty for the plant of having activated immune signaling in the grain (41). A better understanding of NLR regulation by sRNAs and reversible epigenetic marks could inform new strategies to design NLR promoters that maintain the balance between activation of immunity and yield penalty.

3.3. Similarities to Mammalian Intracellular Immunity

Independent evolution of common NLR features is evident not only across plant lineages but also across kingdoms. In animals, a distant homolog of the NB-ARC domain, NACHT, has been



independently combined with LRRs and different signaling domains: CASPASE ACTIVATION AND RECRUITMENT DOMAIN (CARD), BACILLOVIRUS INHIBITOR OF APOPTOSIS PROTEIN REPEAT (BIR), PROTEIN PYRIN DOMAIN (PYD), DEATH DOMAIN (DD), and DEATH EFFECTOR DOMAIN (DED) (127). Mammalian NLRs serve a similar intracellular surveillance function and can induce cell death upon either pathogen perception or autoimmune response (64). In fungi, the NACHT domain-containing protein Het-E triggers nonself recognition and the induction of cell death, called heterokaryon incompatibility (100). Interestingly, the fungal HeLo domain with a four-helix bundle fold similar to RPW8 in RNLs can insert itself into the membrane, thereby forming a pore to induce cell death (38).

A recent breakthrough in the structural biology of plant NLRs has come from solving the structure of the active oligomeric complex of ZAR1 together with its guardees RKS1 and PBL2 kinase (133). The ZAR1 resistosome is structurally similar to the apoptosome of Apaf-1 (151) and the inflammasome of mammalian NLRs (**Figure 2c**) (104, 126, 133). While the structural similarities between the inflammasome and the resistosome are clear, many questions remain unanswered. First, is the CC of CNLs sufficient to form a pore in membranes and induce cell death, or does it require additional factors? The RNLs and their RPW8 domains, which are sufficient to create a pore in structurally similar fungal HeLo proteins, are clear candidates.


Second, do all sensors form homogeneous resistosomes, or does the composition of resistosomes vary? Is there, for example, an equal ratio of paired helpers and sensors, or does a single sensor initiate complex formation of several helpers (**Figure 2c**)? The latter is the case for mammalian NLRs such as NAIP5 and NLRC4 (126). A single NAIP5 sensor is activated by flagellin perception and, upon conformational change, binds to its helper NLRC4, inducing further NLRC4-NLRC4 oligomer formation (126). The final inflammasome forms an open-ring structure with a single NAIP5 sensor and nine NLRC4 helpers. While the ZAR1 resistosome is homogeneous, we cannot exclude the possibility that other plant NLRs can form heterogeneous protein complexes similar to NAIP5/NLRC4. CCs can form both homo- and hetero-oligomers comprising complex interaction networks (30, 86, 139), which would support the idea of heterogeneous resistosomes (**Figure 2c**). While CNLs often form hetero-oligomers (139), reports of TNL hetero-oligomers are relatively rare and are currently limited to paired NLRs (61, 74). It would be exciting to see if the composition of the paired NLR resistosomes is novel and has both sensors and helpers in equal proportions (**Figure 3c**).

The similarity between plant and mammalian cell death execution goes beyond structural similarities of resistosome to inflammasome. Recent reports of NADase activity of bacterial, mammalian and plant TIR domains (47, 48, 59, 131) suggest that activation of cell death can be mediated by conserved secondary messengers. Structural analyses revealed a conserved substrate binding site in the TIR domain of the mammalian executor of neuronal cell death SARM1 and plant TIRs (59). In line with the suggested common function of the TIR proteins across kingdoms, expression of the TIR domain from SARM1 in *Nicotiana benthamiana* produced a cell death response that was visually indistinguishable from HR (59, 131). The major difference between SARM1 and TNLs was revealed by their genetic requirements. Unlike plant TIRs, SARM1 was capable of inducing plant cell death independent of *EDS1* and *NRG1* (59, 131). The elucidation of the precise signaling cascade that culminates in cell death response can reveal further similarities across kingdoms as well as elucidate the plant-specific roles of *EDS1* and *NRG1*.


Finally, does the solved resistosome structure represent its functionally active state? Research on mammalian NLRs suggests that inflammasome formation can be followed by proteasome-mediated release of the active caspase domain (104). Since it has been widely demonstrated in plants that N-terminal truncations of NLRs to either TIR domains or CCs alone are sufficient to induce cell death (16, 30, 52, 69, 122, 139), it is possible that the active resistosome is modified further in the plant cell before the induction of HR.



a Exploiting effector activity



b Integrated domains



c Naturally diverse specificity




Figure 3

Evolutioner’s guide to engineering novel disease resistance. (a) Exploiting effector activities to engineer novel recognition. Prerequisites: a guardee and its corresponding NLR. The introduction of a known effector substrate site into the guardee or the NLR could yield durable resistance when deploying indispensable effectors. (b) Introducing new plant domains in NLR-ID. Prerequisite: an NLR-ID with relatively recent integration. IDs can be exchanged for plant protein domains that are known effector targets. (c) Harvesting natural diversity (indicated by different colors) to identify and subsequently engineer effector binding residues. Prerequisite: availability of information about sequence diversity over a recent period on direct binder NLRs that deploy variable LRR regions for the recognition of effectors. We hypothesize that residues involved in effector binding are more polymorphic and can therefore be identified through natural variation. These residues can be engineered to modify NLR binding specificity to confer altered or even novel recognition capacities.



What can we learn from common patterns observed in NLR evolution? One of the main practical ramifications of the observation that there are multiple ways to evolve the same function is that new functions can also be engineered in multiple ways. Engineering of these functions can be guided by learning the common principles of NLR biology.

4. CURRENT PROGRESS IN ENGINEERING NLR ALLELES

4.1. NLR Chimeras Define Effector Binding Regions and Allow Transfer of Recognition Between Allelic NLRs

Chimeras of flax NLRs were the first successful attempts in changing effector recognition of NLRs (46). This study analyzed 13 alleles of the flax rust resistance gene *L* (*L*, *L1-L11*, and *LH*) to identify variable regions (46). They subsequently created chimeras of *L2*, *L6*, and *L10*; transformed them into flax; and screened for changes in susceptibility to flax rust infection (46). The region exchanged between the NLRs contained 880 amino acids and included the entire LRR region. Transgenic flax plants expressing those chimeras showed resistance and susceptibility phenotypes correlating with the LRR included; in other words, the gene *L6-L2*, containing the *L6* TIR-NB-ARC combined with the *L2* LRR, displayed the resistance phenotype of the *L2* allele (46). This finding demonstrated that the LRR region can be sufficient to mediate recognition specificity of NLRs that directly interact with the effector they recognize. Since the first LRR swaps in *L*, recognition specificity in several other direct binders has been mapped to the LRR region by either allelic swaps (114, 117) or in vivo interactions (26, 34, 44, 69, 102).

4.2. Introduction of New Effector Substrate Sequences into Plant Proteins

NLRs can directly bind effectors or recognize effector function by monitoring a host protein (i.e., a guardee) (**Figure 1b**). Since our understanding of effector functions is continuously improving, exploiting effector activities to engineer resistance seems like an elegant solution. Kim et al. (68) have used this strategy to engineer the guardee of RPS5 in order to expand recognition to several new effector modifications. RPS5 recognizes the action of the bacterial effector AvrPphB, which acts as a protease on PBS1 and related receptor-like cytoplasmic kinases (RLCKs). Upon cleavage by AvrPphB, these RLCKs undergo a conformational change that is recognized by RPS5 and leads to the activation of immune responses (43). The exchange of the proteolytic site in PBS1 with that of another effector, AvrRpt2, drastically improved resistance to *P. syringae* when expressed in otherwise susceptible *Arabidopsis* genotypes (68). The same strategy has also been used to insert the proteolytic site for the viral protease NIa of *Tobacco etch virus* into PBS1 (68). Although this approach led to cleavage of PBS1 through NIa and to HR in *N. benthamiana*, it led to only partial resistance to the virus (68). This finding suggests that rates of substrate cleavage and strength of signaling need to be sufficient to achieve a robust response and may have different requirements for different effectors. Therefore, in some cases this approach might require additional optimization and protein engineering to yield resistance.

4.3. Introduction of Amino Acid Changes to Modify Effector Recognition Specificity

The discovery that the LRR can serve as an effector binding site and mediate NLR recognition specificity has prompted a number of attempts to randomly mutagenize LRRs to generate novel disease resistance. The NLR Rx provides resistance to *Potato virus X* strains carrying a variant



of its coat protein (CP) with the residues T121 and K127 (CP-TK). CPs with a lysine and an arginine in these respective positions (CP-KR) evade recognition by Rx (14). In one study (49), Rx was randomly mutated by means of error-prone polymerase chain reaction, and thousands of variants were screened for gain of recognition of CP-KR. This approach yielded several mutant Rx proteins that provided resistance against strains of *Potato virus x* carrying CP-KR and CP-TK, as well as against another related virus, *Poplar mosaic virus* (49). While this approach expanded the recognition of Rx, it did not yield novel resistance.

Two studies published in 2014 expanded the recognition capacities of the NLR R3a from the wild potato *Solanum demissum* by means of random mutagenesis, gene shuffling, and site-directed mutagenesis (33, 109). The wild-type version of R3a recognizes the *Phytophthora infestans* effector Avr3a^{KI} but not the allelic variant Avr3a^{EM}, which has become a prevalent allele in modern *Phytophthora* species, most likely due to positive selection mediated by evasion of R3a-mediated resistance (5, 20, 21, 130, 146). Segretin et al. (109) identified eight single amino acid substitutions that were able to trigger HR in *N. benthamiana* in response to Avr3a^{EM}. Six of these mutations lie within the LRR region, one in the NB-ARC, and one in the CC. One of these mutations (*K920E*) was also identified in a second study (33). After several rounds of artificial evolution, Chapman et al. (33) identified several mutants that showed a stronger cell death response than the reference. However, these engineered NLRs, termed R3a⁺ and R3a*, were not able to provide enhanced resistance to *P. infestans* strains carrying Avr3a^{EM}.

In a follow-up study, the mutations identified by Segretin and colleagues were transferred to the R3a ortholog I2 from tomato to see whether gain-of-function mutations are transferable between orthologs. I2 confers resistance to *Fusarium oxysporum* f. sp. *lycopersici* through the recognition of the Avr2 effector and has a high sequence similarity to R3a (98, 116). Two of the residues in the R3a⁺ alleles (*I141F* and *N336Y*) are conserved in I2 (60) and were subsequently mutated in I2 in order to expand its recognition capacities (53). However, transfer of these amino acid changes onto I2 led to either loss of recognition or autoactivity (53). To test the hypothesis that these two sites within the homologous NLRs could be hot spots for NLR sensitization, Giannakopoulou et al. (53) mutated both residues to all possible amino acid substitutions. This led to the identification of I2I^{I41N}, which can recognize both Avr3a^{KI} and Avr3a^{EM}, as well as two new Avr2 variants (53). Expression of these NLRs in *N. benthamiana* led to partial resistance to *P. infestans* carrying either variant of Avr3a (53). To date, therefore, attempts to randomly mutagenize NLRs have not led to the generation of novel disease resistance.

5. PROSPECTS AND CHALLENGES IN THE ENGINEERING OF NLRs

5.1. Engineering Guardees for NLR Activation upon a Wide Range of Effector Activities

Exploiting effector activities to engineer novel disease resistance is a promising strategy, as pathogen effectors often evolve to evade NLR binding while retaining their activity. There are several ways in which we can exploit effector activities to the pathogen's detriment. These include the strategy employed by Kim et al. (68) to transfer known effector-targeted proteolytic sites onto a plant protein that is guarded by a plant NLR. We might be able to engineer a guardee whose modification by several effectors can lead to NLR activation (**Figure 3a**). How many different types of effector activities can be recognized by a single guard NLR? The case of RIN4, which is guarded by several NLRs, including RPS2 and RPM1, and is targeted by several effectors with different enzymatic activities suggests that indirect recognition can indeed recognize only one modification, as RPM1 is activated only after phosphorylation of RIN4, while RPS2 recognizes



its proteolytic cleavage (22). The fact that RIN4 is guarded by several NLRs makes it an attractive candidate for the introduction of more effector target sites. The study of effector enzymatic activities will continue to expand the list of potential targets for this approach (22). By choosing effectors with activities that are crucial for pathogen invasion, we can try to create durable resistance. This approach requires comprehensive knowledge of effector activities. Understanding natural effector targets and their activities is therefore crucial for this engineering approach and will greatly enhance our ability to create novel recognition capabilities.

5.2. Engineering NLRs with New Integrated Domains

In addition to modifying guardees, we may be able to engineer NLR clades with IDs. Ideally, we could use existing NLR-IDs as platforms to create fusions with targets identified in effector interactome studies (**Figure 3b**) (101, 136). In practice, the NLR-IDs we observe today have domains that have coevolved for millions of years since the original fusion event. Therefore, this engineering approach will require careful dissection of coevolved regions or identification of a generalist NLR that can still accept new variable domain fusions, such as NLRs from MIC1, identified in Poaceae (10, 121). It will also require a better understanding of NLR-ID activation mechanisms. How does modification of the ID by the effector alter the conformation of the NLR-ID? What is the role of the helper NLR in the pair? Contrasting models have been proposed for different NLR pairs; one suggests that the helper suppresses autoactivity of the sensor NLR, whereas another suggests that the helper mediates initiation of immune signaling (61, 88). Improving our understanding of paired NLRs will facilitate the fine-tuning of immune activation of new NLR-ID fusions.

In the case of the allelic series of *Pik* genes in rice, the integrated heavy-metal-associated (HMA) domain is the most polymorphic region of the NLR (36), strongly suggesting that it is the effector binding interface. This idea was confirmed by crystallization of the *Pik* allele with its corresponding effector (88). A logical next step in utilizing effector binding to an ID is to exchange IDs for other known host domains targeted by effectors (**Figure 3b**). Such an exchange would enable the creation of recognition of effectors whose function remains elusive but whose targets have been identified through their introduction into existing NLR-ID scaffolds (**Figure 3b**). In addition to creating new fusion proteins, protein engineering of existing IDs to alter binding strength between the NLR and effector could be used to improve existing recognition. To this end, knowledge gained from crystal structures of NLRs in association with several allelic effectors, as performed by De La Concepcion et al. (40), has improved our understanding of binding interfaces and which amino acids to alter to increase binding affinity. This information will help guide engineering efforts to extend beyond the random mutagenesis approaches used in the past.

5.3. Exploiting NLRs with Naturally Diverse LRR Binding Specificities

The simplest mode of effector–NLR interaction is direct binding mediated through the variable LRR region, as it involves the fewest genetic components. As a result, direct binder NLRs are potentially the easiest class of NLRs to engineer (**Figure 3c**). However, we currently lack structural and biochemical data for these interactions. Key unanswered questions include the identity of residues within the LRR that determine binding specificity, the typical affinities required for productive interactions, and the structural rearrangements that lead to immune signaling. To sidestep this limitation, we could build upon research with allelic series in the flax *L* gene, discussed above (46), and look to natural allelic diversity and structural modeling to consistently identify binding pockets within the LRRs. By sampling the intraspecific natural diversity of NLRs, we should aim



to determine sets of 10–20 residues within an LRR that are the most variable and, therefore, likely to be involved in effector binding. Such sets of residues will allow targeted engineering efforts to either improve existing binding specificities or derive new ones.

6. CONCLUSIONS AND FUTURE DIRECTIONS

Understanding the evolution of NLRs is crucial in order to determine their functionality, understand the environmental pressures influencing the plant immune receptor repertoire, and engineer new recognition specificities. It is now clear that we cannot apply knowledge of one NLR to another. The phylogenetic placement of NLRs, however, may indicate whether they are more likely to be helpers (e.g., RNLs) or sensors and what type of helpers they depend on (e.g., NRC-dependent clade of sensors). In addition, IDs not only highlight plant proteins that are potentially targeted by pathogens, and therefore monitored by the plant immune system, but also offer unprecedented potential for effector-guided engineering approaches. Nevertheless, it is important to remember that even among genes in one clade there can be functional differences, as is the case for NRG1 performance in Brassicaceae and Solanaceae (25, 71, 102). As sequence databases grow, we will be able to ask increasingly ambitious questions about NLR evolution and find common patterns that were previously concealed by noise.

Engineering of NLRs has proven difficult, and successful reports remain limited. On the basis of our current knowledge of NLRs, there are three main strategies to create novel disease resistance. The recent availability of sequence diversity of NLRs within populations provides the raw material for predictions on effector binding surfaces. For NLR-IDs, it is already apparent that IDs could potentially serve as effector target platforms. We now require an understanding of the timing of fusions and subsequent coevolution with canonical NLR domains. This information will inform the choice of a scaffold NLR that could be used for the exchange of IDs. For instance, NLR-IDs from the evolutionary youngest clade, MIC1, likely possess the greatest potential for successful engineering approaches. In the near future, lessons learned from NLR evolution and advances in synthetic and structural biology will enable us to create highly coveted designer NLRs.

SUMMARY POINTS

1. Subfunctionalization of nucleotide-binding leucine-rich repeat receptors (NLRs) into sensors and helpers arose independently multiple times across different species.
2. Helpers that evolved from NLR pairs are evolutionarily younger than coiled-coil domain NLR (RNL) helpers.
3. Subfunctionalization in sensor and helper signaling components of NLRs is reminiscent of the receptor and coreceptor relationship of pattern recognition receptors and might be a general evolutionary theme.
4. While predicting the mode of action of NLRs from sequence alone is challenging, phylogenetic placement in clades can help predict their genetic requirements.
5. Engineering of NLRs has been difficult, with only moderate success thus far.
6. Understanding the evolution of NLRs can inform engineering efforts and lead to designer NLRs with novel recognition specificity.



FUTURE ISSUES

1. Why do plants lose certain NLR clades but expand others?
2. Are evolutionary patterns conserved between intraspecific and interspecific scales?
3. How do human and pathogen evolutionary pressures affect immune gene evolution? Is this consistent across systems and between domesticated and wild systems?
4. On what timescale and by what means is the regulation of NLRs evolving in response to varying fitness costs with changing environments? Is this occurring at a faster or slower rate than sequence evolution within the NLR?
5. Why do NLRs subfunctionalize into helpers and sensors? How does this relationship work, and how is this complex activated by an effector?
6. What is the functional link between ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1), PHYTOALEXIN DEFICIENT 4 (PAD4), and RNL helpers? What are the exact molecular events leading to the hypersensitive response (HR), and what is its function in plant immunity?
7. What is the function of integrated domains (IDs)? Are they merely pathogen bait, or do they also fulfill a function by, for instance, targeting the NLR to the correct subcellular compartment?
8. Which NLR-IDs can serve as platforms for new fusion proteins? Which guardees can serve as platforms for effector target domains?

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank Dr. Paul Bailey for generously providing the NLR tree included as a figure in this review. We thank Erin L. Baggs and Dr. Daniil Prigozhin for insightful discussions of the concepts presented in this review and for critical reading of the manuscript. We also thank Pierre Joubert for providing his comments on the review.

LITERATURE CITED

1. Aarts N, Metz M, Holub E, Staskawicz BJ, Daniels MJ, Parker JE. 1998. Different requirements for EDS1 and NDR1 by disease resistance genes define at least two *R* gene-mediated signaling pathways in *Arabidopsis*. *PNAS* 95:10306–11
2. Adachi H, Contreras MP, Harant A, Wu C-H, Derevnina L, et al. 2019. An N-terminal motif in NLR immune receptors is functionally conserved across distantly related plant species. *eLife* 8:e49956
3. Alcázar R, von Reth M, Bautor J, Chae E, Weigel D, et al. 2014. Analysis of a plant complex resistance gene locus underlying immune-related hybrid incompatibility and its occurrence in nature. *PLOS Genet.* 10:e1004848
4. Andersen EJ, Nepal MP. 2019. Diversification of disease resistance receptors by integrated domain fusions in wheat and its progenitors. bioRxiv 695148. <https://doi.org/10.1101/695148>



5. Armstrong MR, Whisson SC, Pritchard L, Bos JIB, Venter E, et al. 2005. An ancestral oomycete locus contains late blight avirulence gene *Avr3a*, encoding a protein that is recognized in the host cytoplasm. *PNAS* 102:7766–71
6. Ashfield T, Redditt T, Russell A, Kessens R, Rodibaugh N, et al. 2014. Evolutionary relationship of disease resistance genes in soybean and *Arabidopsis* specific for the *Pseudomonas syringae* effectors AvrB and AvrRpm1. *Plant Physiol.* 166:235–51
7. Axtell MJ, Staskawicz BJ. 2003. Initiation of RPS2-specified disease resistance in *Arabidopsis* is coupled to the AvrRpt2-directed elimination of RIN4. *Cell* 112:369–77
8. Baggs E, Dagdas G, Krasileva KV. 2017. NLR diversity, helpers and integrated domains: making sense of the NLR IDentity. *Curr. Opin. Plant Biol.* 38:59–67
9. Baggs EL, Thanki AS, O’Grady R, Schudoma C, Haerty W, Krasileva KV. 2019. Convergent gene loss in aquatic plants predicts new components of plant immunity and drought response. bioRxiv 572560. <https://doi.org/10.1101/572560>
10. Bailey PC, Schudoma C, Jackson W, Baggs E, Dagdas G, et al. 2018. Dominant integration locus drives continuous diversification of plant immune receptors with exogenous domain fusions. *Genome Biol.* 19:23
11. Barragan CA, Wu R, Kim S-T, Xi W, Habring A, et al. 2019. RPW8/HR repeats control NLR activation in *Arabidopsis thaliana*. *PLoS Genet.* 15:e1008313
12. Bastedo DP, Khan M, Martel A, Seto D, Kireeva I, et al. 2019. Perturbations of the ZED1 pseudokinase activate plant immunity. *PLoS Pathog.* 15:e1007900
13. Belkhadir Y, Nimchuk Z, Hubert DA, Mackey D, Dangl JL. 2004. *Arabidopsis* RIN4 negatively regulates disease resistance mediated by RPS2 and RPM1 downstream or independent of the NDR1 signal modulator and is not required for the virulence functions of bacterial type III effectors AvrRpt2 or AvrRpm1. *Plant Cell* 16:2822–35
14. Bendahmane A, Köhm BA, Dedi C, Baulcombe DC. 1995. The coat protein of potato virus X is a strain-specific elicitor of Rx1-mediated virus resistance in potato. *Plant J.* 8:933–41
15. Bernoux M, Burdett H, Williams SJ, Zhang X, Chen C, et al. 2016. Comparative analysis of the flax immune receptors L6 and L7 suggests an equilibrium-based switch activation model. *Plant Cell* 28:146–59
16. Bernoux M, Ve T, Williams S, Warren C, Hatters D, et al. 2011. Structural and functional analysis of a plant resistance protein TIR domain reveals interfaces for self-association, signaling, and autoregulation. *Cell Host Microbe* 9:200–11
17. Bittner-Eddy PD, Crute IR, Holub EB, Beynon JL. 2000. *RPP13* is a simple locus in *Arabidopsis thaliana* for alleles that specify downy mildew resistance to different avirulence determinants in *Peronospora parasitica*. *Plant J.* 21:177–88
18. Boller T, Felix G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu. Rev. Plant Biol.* 60:379–406
19. Bombliès K, Lempe J, Epple P, Warthmann N, Lanz C, et al. 2007. Autoimmune response as a mechanism for a Dobzhansky-Muller-type incompatibility syndrome in plants. *PLoS Biol.* 5:e236
20. Bos JIB, Chaparro-García A, Quesada-Ocampo LM, McSpadden Gardener BB, Kamoun S. 2009. Distinct amino acids of the *Phytophthora infestans* effector AVR3a condition activation of R3a hypersensitivity and suppression of cell death. *Mol. Plant-Microbe Interact.* 22:269–81
21. Bos JIB, Kanneganti T-D, Young C, Cakir C, Huitema E, et al. 2006. The C-terminal half of *Phytophthora infestans* RXLR effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and suppress INF1-induced cell death in *Nicotiana benthamiana*. *Plant J.* 48:165–76
22. Büttner D. 2016. Behind the lines—actions of bacterial type III effector proteins in plant cells. *FEMS Microbiol. Rev.* 40:894–937
23. Cambiagno DA, Nota F, Zavallo D, Rius S, Casati P, et al. 2018. Immune receptor genes and pericentromeric transposons as targets of common epigenetic regulatory elements. *Plant J.* 96:1178–90
24. Carter ME, Helm M, Chapman AVE, Wan E, Restrepo Sierra AM, et al. 2019. Convergent evolution of effector protease recognition by *Arabidopsis* and barley. *Mol. Plant-Microbe Interact.* 32:550–65
25. Castel B, Ngou P-M, Cevik V, Redkar A, Kim D-S, et al. 2019. Diverse NLR immune receptors activate defence via the RPW8-NLR NRG1. *New Phytol.* 222:966–80



26. Catanzariti A-M, Dodds PN, Ve T, Kobe B, Ellis JG, Staskawicz BJ. 2010. The AvrM effector from flax rust has a structured C-terminal domain and interacts directly with the M resistance protein. *Mol. Plant-Microbe Interact.* 23:49–57
27. Century KS, Shapiro AD, Repetti PP, Dahlbeck D, Holub E, Staskawicz BJ. 1997. NDR1, a pathogen-induced component required for *Arabidopsis* disease resistance. *Science* 278:1963–65
28. Césari S, Bernoux M, Moncuquet P, Kroj T, Dodds PN. 2014. A novel conserved mechanism for plant NLR protein pairs: the “integrated decoy” hypothesis. *Front. Plant Sci.* 5:606
29. Césari S, Kanzaki H, Fujiwara T, Bernoux M, Chalvon V, et al. 2014. The NB-LRR proteins RGA4 and RGA5 interact functionally and physically to confer disease resistance. *EMBO J.* 33:1941–59
30. Césari S, Moore J, Chen C, Webb D, Periyannan S, et al. 2016. Cytosolic activation of cell death and stem rust resistance by cereal MLA-family CC-NLR proteins. *PNAS* 113:10204–9
31. Chae E, Bomblies K, Kim S-T, Karelina D, Zaidem M, et al. 2014. Species-wide genetic incompatibility analysis identifies immune genes as hot spots of deleterious epistasis. *Cell* 159:1341–51
32. Chae E, Tran DTN, Weigel D. 2016. Cooperation and conflict in the plant immune system. *PLOS Pathog.* 12:e1005452
33. Chapman S, Stevens LJ, Boevink PC, Engelhardt S, Alexander CJ, et al. 2014. Detection of the virulent form of AVR3a from *Phytophthora infestans* following artificial evolution of potato resistance gene *R3a*. *PLOS ONE* 9:e110158
34. Chen J, Upadhyaya NM, Ortiz D, Sperschneider J, Li F, et al. 2017. Loss of by somatic exchange in stem rust leads to virulence for resistance in wheat. *Science* 358:1607–10
35. Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nürnberger T, et al. 2007. A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* 448:497–500
36. Costanzo S, Jia Y. 2010. Sequence variation at the rice blast resistance gene *Pi-km* locus: implications for the development of allele specific markers. *Plant Sci.* 178:523–30
37. Couto D, Zipfel C. 2016. Regulation of pattern recognition receptor signalling in plants. *Nat. Rev. Immunol.* 16:537–52
38. Daskalov A, Habenstein B, Sabaté R, Berbon M, Martinez D, et al. 2016. Identification of a novel cell death-inducing domain reveals that fungal amyloid-controlled programmed cell death is related to necroptosis. *PNAS* 113:2720–25
39. Day B, Dahlbeck D, Staskawicz BJ. 2006. NDR1 interaction with RIN4 mediates the differential activation of multiple disease resistance pathways in *Arabidopsis*. *Plant Cell* 18:2782–91
40. **De La Concepcion JC, Franceschetti M, Maqbool A, Saitoh H, Terauchi R, et al. 2018. Polymorphic residues in rice NLRs expand binding and response to effectors of the blast pathogen. *Nat. Plants* 4:576–85**
41. Deng Y, Zhai K, Xie Z, Yang D, Zhu X, et al. 2017. Epigenetic regulation of antagonistic receptors confers rice blast resistance with yield balance. *Science* 355:962–65
42. Deslandes L, Olivier J, Peeters N, Feng DX, Khounlotham M, et al. 2003. Physical interaction between RRS1-R, a protein conferring resistance to bacterial wilt, and PopP2, a type III effector targeted to the plant nucleus. *PNAS* 100:8024–29
43. DeYoung BJ, Qi D, Kim S-H, Burke TP, Innes RW. 2012. Activation of a plant nucleotide binding–leucine rich repeat disease resistance protein by a modified self protein. *Cell. Microbiol.* 14:1071–84
44. Dodds PN, Lawrence GJ, Catanzariti A-M, Teh T, Wang C-IA, et al. 2006. Direct protein interaction underlies gene-for-gene specificity and coevolution of the flax resistance genes and flax rust avirulence genes. *PNAS* 103:8888–93
45. Dodds PN, Rathjen JP. 2010. Plant immunity: towards an integrated view of plant–pathogen interactions. *Nat. Rev. Genet.* 11:539–48
46. Ellis JG, Lawrence GJ, Luck JE, Dodds PN. 1999. Identification of regions in alleles of the flax rust resistance gene *L* that determine differences in gene-for-gene specificity. *Plant Cell* 11:495–506
47. Essuman K, Summers DW, Sasaki Y, Mao X, DiAntonio A, Milbrandt J. 2017. The SARM1 Toll/interleukin-1 receptor domain possesses intrinsic NAD cleavage activity that promotes pathological axonal degeneration. *Neuron* 93:1334–43

40. Provides genetic and biochemical data showing the expanded range of effector recognition by NLR-HMA.



48. Essuman K, Summers DW, Sasaki Y, Mao X, Yim AKY, et al. 2018. TIR domain proteins are an ancient family of NAD-consuming enzymes. *Curr. Biol.* 28:421–30
49. Farnham G, Baulcombe DC. 2006. Artificial evolution extends the spectrum of viruses that are targeted by a disease-resistance gene from potato. *PNAS* 103:18828–33
50. Feys BJ, Moisan LJ, Newman MA, Parker JE. 2001. Direct interaction between the *Arabidopsis* disease resistance signaling proteins, EDS1 and PAD4. *EMBO J.* 20:5400–11
51. Flor HH. 1955. Host-parasite interaction in flax rust—its genetics and other implications. *Phytopathology* 45:680–85
52. Frost D, Way H, Howles P, Luck J, Manners J, et al. 2004. Tobacco transgenic for the flax rust resistance gene *L* expresses allele-specific activation of defense responses. *Mol. Plant-Microbe Interact.* 17:224–32
53. Giannakopoulou A, Steele JFC, Segretin ME, Bozkurt TO, Zhou J, et al. 2015. Tomato I2 immune receptor can be engineered to confer partial resistance to the oomycete *Phytophthora infestans* in addition to the fungus *Fusarium oxysporum*. *Mol. Plant-Microbe Interact.* 28:1316–29
54. González VM, Müller S, Baulcombe D, Puigdomènech P. 2015. Evolution of NBS-LRR gene copies among dicot plants and its regulation by members of the miR482/2118 superfamily of miRNAs. *Mol. Plant* 8:329–31
55. Gu L, Si W, Zhao L, Yang S, Zhang X. 2015. Dynamic evolution of *NBS-LRR* genes in bread wheat and its progenitors. *Mol. Genet. Genom.* 290:727–38
56. Guo Y-L, Fitz J, Schneeberger K, Ossowski S, Cao J, Weigel D. 2011. Genome-wide comparison of nucleotide-binding site leucine-rich repeat-encoding genes in *Arabidopsis*. *Plant Physiol.* 157:757–69
57. Gust AA, Pruitt R, Nürnberger T. 2017. Sensing danger: key to activating plant immunity. *Trends Plant Sci.* 22:779–91
58. He X-F, Fang Y-Y, Feng L, Guo H-S. 2008. Characterization of conserved and novel microRNAs and their targets, including a TuMV-induced TIR-NBS-LRR class *R* gene-derived novel miRNA in *Brassica*. *FEBS Lett.* 582:2445–52
59. Horsefield S, Burdett H, Zhang X, Manik MK, Shi Y, et al. 2019. NAD⁺ cleavage activity by animal and plant TIR domains in cell death pathways. *Science* 365:793–99
60. Huang S, van der Vossen EAG, Kuang H, Vleeshouwers VGAA, Zhang N, et al. 2005. Comparative genomics enabled the isolation of the *R3a* late blight resistance gene in potato. *Plant J.* 42:251–61
61. Huh SU, Cevik V, Ding P, Duxbury Z, Ma Y, et al. 2017. Protein–protein interactions in the RPS4/RRS1 immune receptor complex. *PLOS Pathog.* 13:e1006376
62. Jia Y, McAdams SA, Bryan GT, Hershey HP, Valent B. 2000. Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. *EMBO J.* 19:4004–14
63. Jia Y, Yuan Y, Zhang Y, Yang S, Zhang X. 2015. Extreme expansion of NBS-encoding genes in Rosaceae. *BMC Genet.* 16:48
64. Jones JDG, Vance RE, Dangl JL. 2016. Intracellular innate immune surveillance devices in plants and animals. *Science* 354:aaf6395
65. Jubic LM, Saile S, Furzer OJ, El Kasm F, Dangl JL. 2019. Help wanted: helper NLRs and plant immune responses. *Curr. Opin. Plant Biol.* 50:82–94
66. Karasov TL, Kniskern JM, Gao L, DeYoung BJ, Ding J, et al. 2014. The long-term maintenance of a resistance polymorphism through diffuse interactions. *Nature* 512:436–40
67. Kim S, Park J, Yeom S-I, Kim Y-M, Seo E, et al. 2017. New reference genome sequences of hot pepper reveal the massive evolution of plant disease-resistance genes by retroduplication. *Genome Biol.* 18:210
68. **Kim SH, Qi D, Ashfield T, Helm M, Innes RW. 2016. Using decoys to expand the recognition specificity of a plant disease resistance protein. *Science* 351:684–87**
69. Krasileva KV, Dahlbeck D, Staskawicz BJ. 2010. Activation of an *Arabidopsis* resistance protein is specified by the in planta association of its leucine-rich repeat domain with the cognate oomycete effector. *Plant Cell* 22:2444–58
70. Kroj T, Chanclud E, Michel-Romiti C, Grand X, Morel J-B. 2016. Integration of decoy domains derived from protein targets of pathogen effectors into plant immune receptors is widespread. *New Phytol.* 210:618–26

68. Introduces an exogenous protease cleavage site into a guard cell to engineer resistance to a new pathogen.



71. Lapin D, Kovacova V, Sun X, Dongus JA, Bhandari DD, et al. 2019. A coevolved EDS1-SAG101-NRG1 module mediates cell death signaling by TIR-domain immune receptors. *Plant Cell* 31:2430–55
72. Leister D. 2004. Tandem and segmental gene duplication and recombination in the evolution of plant disease resistance genes. *Trends Genet.* 20:116–22
73. Lewis JD, Lee AH-Y, Hassan JA, Wan J, Hurley B, et al. 2013. The *Arabidopsis* ZED1 pseudokinase is required for ZAR1-mediated immunity induced by the *Pseudomonas syringae* type III effector HopZ1a. *PNAS* 110:18722–27
74. Liang W, van Wersch S, Tong M, Li X. 2019. TIR-NB-LRR immune receptor SOC3 pairs with truncated TIR-NB protein CHS1 or TN2 to monitor the homeostasis of E3 ligase SAUL1. *New Phytol.* 221:2054–66
75. Liang X, Zhou J-M. 2018. Receptor-like cytoplasmic kinases: central players in plant receptor kinase-mediated signaling. *Annu. Rev. Plant Biol.* 69:267–99
76. Li F, Pignatta D, Bendix C, Brunkard JO, Cohn MM, et al. 2012. MicroRNA regulation of plant innate immune receptors. *PNAS* 109:1790–95
77. Li L, Habring A, Wang K, Weigel D. 2019. Oligomerization of NLR immune receptor RPP7 triggered by atypical resistance protein RPW8/HR as ligand. bioRxiv 682807. <https://doi.org/10.1101/682807>
78. Lin X, Zhang Y, Kuang H, Chen J. 2013. Frequent loss of lineages and deficient duplications accounted for low copy number of disease resistance genes in Cucurbitaceae. *BMC Genom.* 14:335
79. Liu C, Cui D, Zhao J, Liu N, Wang B, et al. 2019. Two *Arabidopsis* receptor-like cytoplasmic kinases SZE1 and SZE2 associate with the ZAR1-ZED1 complex and are required for effector-triggered immunity. *Mol. Plant* 12:967–83
80. Liu J, Cheng X, Liu D, Xu W, Wise R, Shen Q-H. 2014. The miR9863 family regulates distinct *Mla* alleles in barley to attenuate NLR receptor-triggered disease resistance and cell-death signaling. *PLoS Genet.* 10:e1004755
81. Li Y, Zhong Y, Huang K, Cheng Z-M. 2016. Genomewide analysis of NBS-encoding genes in kiwi fruit (*Actinidia chinensis*). *J. Genet.* 95:997–1001
82. Luo S, Zhang Y, Hu Q, Chen J, Li K, et al. 2012. Dynamic nucleotide-binding site and leucine-rich repeat-encoding genes in the grass family. *Plant Physiol.* 159:197–210
83. Lu X, Kracher B, Saur IML, Bauer S, Ellwood SR, et al. 2016. Allelic barley MLA immune receptors recognize sequence-unrelated avirulence effectors of the powdery mildew pathogen. *PNAS* 113:E6486–95
84. Mackey D, Belkhadir Y, Alonso JM, Ecker JR, Dangl JL. 2003. *Arabidopsis* RIN4 is a target of the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. *Cell* 112:379–89
85. Mackey D, Holt BF 3rd, Wiig A, Dangl JL. 2002. RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* 108:743–54
86. Maekawa T, Cheng W, Spiridon LN, Töller A, Lukasik E, et al. 2011. Coiled-coil domain-dependent homodimerization of intracellular barley immune receptors defines a minimal functional module for triggering cell death. *Cell Host Microbe* 9:187–99
87. Mago R, Zhang P, Vautrin S, Šimková H, Bansal U, et al. 2015. The wheat *Sr50* gene reveals rich diversity at a cereal disease resistance locus. *Nat Plants* 1:15186
88. Maqbool A, Saitoh H, Franceschetti M, Stevenson CEM, Uemura A, et al. 2015. Structural basis of pathogen recognition by an integrated HMA domain in a plant NLR immune receptor. *eLife* 4:e08709
89. Ma Y, Guo H, Hu L, Martinez PP, Moschou PN, et al. 2018. Distinct modes of derepression of an immune receptor complex by two different bacterial effectors. *PNAS* 115:10218–27
90. Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. 2003. Genome-wide analysis of NBS-LRR-encoding genes in *Arabidopsis*. *Plant Cell* 15:809–34
91. Michelmore RW, Meyers BC. 1998. Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. *Genome Res.* 8:1113–30
92. Mondragon-Palomino M, Gaut BS. 2005. Gene conversion and the evolution of three leucine-rich repeat gene families in *Arabidopsis thaliana*. *Mol. Biol. Evol.* 22:2444–56
93. Nam KH, Li J. 2002. BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signaling. *Cell* 110:203–12



94. Nandety RS, Caplan JL, Cavanaugh K, Perroud B, Wroblewski T, et al. 2013. The role of TIR-NBS and TIR-X proteins in plant basal defense responses. *Plant Physiol.* 162:1459–72
95. Narusaka M, Kubo Y, Shiraishi T, Iwabuchi M, Narusaka Y. 2009. A dual resistance gene system prevents infection by three distinct pathogens. *Plant Signal. Behav.* 4:954–55
96. Nishimura MT, Anderson RG, Cherkis KA, Law TF, Liu QL, et al. 2017. TIR-only protein RBA1 recognizes a pathogen effector to regulate cell death in *Arabidopsis*. *PNAS* 114:E2053–62
97. Nuthikattu S, McCue AD, Panda K, Fultz D, DeFraia C, et al. 2013. The initiation of epigenetic silencing of active transposable elements is triggered by RDR6 and 21–22 nucleotide small interfering RNAs. *Plant Physiol.* 162:116–31
98. Ori N, Eshed Y, Paran I, Presting G, Aviv D, et al. 1997. The I2C family from the wilt disease resistance locus I2 belongs to the nucleotide binding, leucine-rich repeat superfamily of plant resistance genes. *Plant Cell* 9:521–32
99. Ouyang S, Park G, Atamian HS, Han CS, Stajich JE, et al. 2014. MicroRNAs suppress NB domain genes in tomato that confer resistance to *Fusarium oxysporum*. *PLOS Pathog.* 10:e1004464
100. Paoletti M, Clavé C. 2007. The fungus-specific HET domain mediates programmed cell death in *Podospora anserina*. *Eukaryot. Cell* 6:2001–8
101. Petre B, Saunders DGO, Sklenar J, Lorrain C, Krasileva KV, et al. 2016. Heterologous expression screens in *Nicotiana benthamiana* identify a candidate effector of the wheat yellow rust pathogen that associates with processing bodies. *PLOS ONE* 11:e0149035
102. Qi T, Seong K, Thomazella DPT, Kim JR, Pham J, et al. 2018. NRG1 functions downstream of EDS1 to regulate TIR-NLR-mediated plant immunity in *Nicotiana benthamiana*. *PNAS* 115:E10979–87
103. Rodin SN, Riggs AD. 2003. Epigenetic silencing may aid evolution by gene duplication. *J. Mol. Evol.* 56:718–29
104. Sandstrom A, Mitchell PS, Goers L, Mu EW, Lesser CF, Vance RE. 2019. Functional degradation: a mechanism of NLRP1 inflammasome activation by diverse pathogen enzymes. *Science* 364:eaau1330
105. Sarris PF, Cevik V, Dagdas G, Jones JDG, Krasileva KV. 2016. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. *BMC Biol.* 14:8
106. Sarris PF, Duxbury Z, Huh SU, Ma Y, Segonzac C, et al. 2015. A plant immune receptor detects pathogen effectors that target WRKY transcription factors. *Cell* 161:1089–100
107. Saucet SB, Ma Y, Sarris PF, Furzer OJ, Sohn KH, Jones JDG. 2015. Two linked pairs of *Arabidopsis* TNL resistance genes independently confer recognition of bacterial effector AvrRps4. *Nat. Commun.* 6:6338
108. Schultink A, Qi T, Bally J, Staskawicz B. 2019. Using forward genetics in *Nicotiana benthamiana* to uncover the immune signaling pathway mediating recognition of the *Xanthomonas perforans* effector XopJ4. *New Phytol.* 221:1001–9
109. Segretin ME, Pais M, Franceschetti M, Chaparro-Garcia A, Bos JIB, et al. 2014. Single amino acid mutations in the potato immune receptor R3a expand response to *Phytophthora* effectors. *Mol. Plant-Microbe Interact.* 27:624–37
110. Seo E, Kim T, Park JH, Yeom S-I, Kim S, et al. 2018. Genome-wide comparative analysis in Solanaceae species reveals evolution of microRNAs targeting defense genes in *Capsicum* spp. *DNA Res.* 25:561–75
111. Serra H, Choi K, Zhao X, Blackwell AR, Kim J, Henderson IR. 2018. Interhomolog polymorphism shapes meiotic crossover within the *Arabidopsis* *RAC1* and *RPP13* disease resistance genes. *PLOS Genet.* 14:e1007843
112. Seto D, Koulina N, Lo T, Menna A, Guttman DS, Desveaux D. 2017. Expanded type III effector recognition by the ZARI NLR protein using ZED1-related kinases. *Nat. Plants* 3:17027
113. Shao Z-Q, Xue J-Y, Wu P, Zhang Y-M, Wu Y, et al. 2016. Large-scale analyses of angiosperm nucleotide-binding site-leucine-rich repeat genes reveal three anciently diverged classes with distinct evolutionary patterns. *Plant Physiol.* 170:2095–109
114. Shen Q-H, Zhou F, Bieri S, Haizel T, Shirasu K, Schulze-Lefert P. 2003. Recognition specificity and RAR1/SGT1 dependence in barley *Mla* disease resistance genes to the powdery mildew fungus. *Plant Cell* 15:732–44

113. Analyzes 22 angiosperms grouped in NLRs in three major clades and reconstructs their evolutionary history.



115. Shivaprasad PV, Chen H-M, Patel K, Bond DM, Santos BACM, Baulcombe DC. 2012. A microRNA superfamily regulates nucleotide binding site–leucine-rich repeats and other mRNAs. *Plant Cell* 24:859–74
116. Simons G, Groenendijk J, Wijbrandi J, Reijans M, Groenen J, et al. 1998. Dissection of the *Fusarium I2* gene cluster in tomato reveals six homologs and one active gene copy. *Plant Cell* 10:1055–68
117. Slootweg E, Koropacka K, Roosien J, Dees R, Overmars H, et al. 2017. Sequence exchange between homologous NB-LRR genes converts virus resistance into nematode resistance, and vice versa. *Plant Physiol.* 175:498–510
118. Song W, Wang B, Li X, Wei J, Chen L, et al. 2015. Identification of immune related LRR-containing genes in maize (*Zea mays* L.) by genome-wide sequence analysis. *Int. J. Genom. Proteom.* 2015:231358
119. Stam R, Scheikl D, Tellier A. 2016. Pooled enrichment sequencing identifies diversity and evolutionary pressures at NLR resistance genes within a wild tomato population. *Genome Biol. Evol.* 8:1501–15
120. Stam R, Silva-Arias GA, Tellier A. 2019. Subsets of NLR genes show differential signatures of adaptation during colonization of new habitats. *New Phytol.* 224:367–79
121. Stein JC, Yu Y, Copetti D, Zwickl DJ, Zhang L, et al. 2018. Genomes of 13 domesticated and wild rice relatives highlight genetic conservation, turnover and innovation across the genus *Oryza*. *Nat. Genet.* 50:285–96
122. Swiderski MR, Birker D, Jones JDG. 2009. The TIR domain of TIR-NB-LRR resistance proteins is a signaling domain involved in cell death induction. *Mol. Plant-Microbe Interact.* 22:157–65
123. Takken FLW, Albrecht M, Tameling WI. 2006. Resistance proteins: molecular switches of plant defence. *Curr. Opin. Plant Biol.* 9:383–90
124. Takken FLW, Govers A. 2012. How to build a pathogen detector: structural basis of NB-LRR function. *Curr. Opin. Plant Biol.* 15:375–84
125. Tameling WIL, Elzinga SDJ, Darmin PS, Vossen JH, Takken FLW, et al. 2002. The tomato *R* gene products I-2 and MI-1 are functional ATP binding proteins with ATPase activity. *Plant Cell* 14:2929–39
126. Tenthorey JL, Haloupek N, López-Blanco JR, Grob P, Adamson E, et al. 2017. The structural basis of flagellin detection by NAIP5: a strategy to limit pathogen immune evasion. *Science* 358:888–93
127. Urbach JM, Ausubel FM. 2017. The NBS-LRR architectures of plant R-proteins and metazoan NLRs evolved in independent events. *PNAS* 114:1063–68
128. Van de Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, et al. 2019. A species-wide inventory of NLR genes and alleles in *Arabidopsis thaliana*. *Cell* 178:1260–72
129. van der Hoorn RAL, Kamoun S. 2008. From guard to decoy: a new model for perception of plant pathogen effectors. *Plant Cell.* 20:2009–17
130. Vleeshouwers VGAA, Raffaele S, Vossen JH, Champouret N, Oliva R, et al. 2011. Understanding and exploiting late blight resistance in the age of effectors. *Annu. Rev. Phytopathol.* 49:507–31
131. Wan L, Essuman K, Anderson RG, Sasaki Y, Monteiro F, et al. 2019. TIR domains of plant immune receptors are NAD⁺-cleaving enzymes that promote cell death. *Science.* 365:799–803
132. Wang G, Roux B, Feng F, Guy E, Li L, et al. 2015. The decoy substrate of a pathogen effector and a pseudokinase specify pathogen-induced modified-self recognition and immunity in plants. *Cell Host Microbe* 18:285–95
133. Wang J, Hu M, Wang J, Qi J, Han Z, et al. 2019. Reconstitution and structure of a plant NLR resistosome conferring immunity. *Science* 364:eaav5870
134. Wang J, Wang J, Hu M, Wu S, Qi J, et al. 2019. Ligand-triggered allosteric ADP release primes a plant NLR complex. *Science* 364:eaav5868
135. Wang W, Devoto A, Turner JG, Xiao S. 2007. Expression of the membrane-associated resistance protein RPW8 enhances basal defense against biotrophic pathogens. *Mol. Plant-Microbe Interact.* 20:966–76
136. Weßling R, Epple P, Altmann S, He Y, Yang L, et al. 2014. Convergent targeting of a common host protein network by pathogen effectors from three kingdoms of life. *Cell Host Microbe* 16:364–75
137. Williams SJ, Sornaraj P, deCourcy-Ireland E, Menz RI, Kobe B, et al. 2011. An autoactive mutant of the *M* flax rust resistance protein has a preference for binding ATP, whereas wild-type *M* protein binds ADP. *Mol. Plant-Microbe Interact.* 24:897–906
133. Reveals the wheel-shaped pentameric resistosome structure of the active ZAR1 complex.
134. Provides biochemical and structural data for intramolecular activation of ZAR1 upon effector perception.



139. Shows that CC domains from *Arabidopsis* NLRs form a network of homo- and heteromeric interactions.

149. Analyzes 70 plants, showing the rapid birth of new miRNAs from recently duplicated NLRs.

138. Williams SJ, Yin L, Foley G, Casey LW, Outram MA, et al. 2016. Structure and function of the TIR domain from the grape NLR protein RPV1. *Front. Plant Sci.* 7:1850
139. Wróblewski T, Spiridon L, Martin EC, Petrescu A-J, Cavanaugh K, et al. 2018. Genome-wide functional analyses of plant coiled-coil NLR-type pathogen receptors reveal essential roles of their N-terminal domain in oligomerization, networking, and immunity. *PLOS Biol.* 16:e2005821
140. Wu C-H, Abd-El-Haliem A, Bozkurt TO, Belhaj K, Terauchi R, et al. 2017. NLR network mediates immunity to diverse plant pathogens. *PNAS* 114:8113–18
141. Wu C-H, Belhaj K, Bozkurt TO, Birk MS, Kamoun S. 2016. Helper NLR proteins NRC2a/b and NRC3 but not NRC1 are required for Pto-mediated cell death and resistance in *Nicotiana benthamiana*. *New Phytol.* 209:1344–52
142. Wu C-H, Kamoun S. 2019. Tomato Prf requires NLR helpers NRC2 and NRC3 to confer resistance against the bacterial speck pathogen *Pseudomonas syringae* pv. *tomato*. bioRxiv 595744. <https://doi.org/10.1101/595744>
143. Wu Z, Li M, Dong OX, Xia S, Liang W, et al. 2019. Differential regulation of TNL-mediated immune signaling by redundant helper CNLs. *New Phytol.* 222:938–53
144. Xiao S, Ellwood S, Calis O, Patrick E, Li T, et al. 2001. Broad-spectrum mildew resistance in *Arabidopsis thaliana* mediated by RPW8. *Science* 291:118–20
145. Xia R, Xu J, Arikis S, Meyers BC. 2015. Extensive families of miRNAs and PHAS loci in Norway spruce demonstrate the origins of complex phasiRNA networks in seed plants. *Mol. Biol. Evol.* 32:2905–18
146. Yoshida K, Schuenemann VJ, Cano LM, Pais M, Mishra B, et al. 2013. The rise and fall of the *Phytophthora infestans* lineage that triggered the Irish potato famine. *eLife* 2:e00731
147. Zhai J, Jeong D-H, De Paoli E, Park S, Rosen BD, et al. 2011. MicroRNAs as master regulators of the plant NB-LRR defense gene family via the production of phased, *trans*-acting siRNAs. *Genes Dev.* 25:2540–53
148. Zhang X, Bernoux M, Bentham AR, Newman TE, Ve T, et al. 2017. Multiple functional self-association interfaces in plant TIR domains. *PNAS* 114:E2046–52
149. Zhang Y, Xia R, Kuang H, Meyers BC. 2016. The diversification of plant NBS-LRR defense genes directs the evolution of microRNAs that target them. *Mol. Biol. Evol.* 33:2692–705
150. Zhang Z, Wu Y, Gao M, Zhang J, Kong Q, et al. 2012. Disruption of PAMP-induced MAP kinase cascade by a *Pseudomonas syringae* effector activates plant immunity mediated by the NB-LRR protein SUMM2. *Cell Host Microbe* 11:253–63
151. Zhou M, Li Y, Hu Q, Bai X, Huang W, et al. 2015. Atomic structure of the apoptosome: mechanism of cytochrome *c*- and dATP-mediated activation of Apaf-1. *Genes Dev.* 29:2349–61

